



GI For Kids

Pediatric Gastroenterology and Nutrition Services

Excellent Care Every Time

1975 Town Center Blvd · Knoxville, TN 37922

Phone (865) 546-3998 · Fax (865) 546-1123 · www.giforkids.com

Colonoscopy Prep Ages 6-10 Turkey Creek Medical Center – Tennova

Prep Date:

Procedure Date:

It is very important to make sure that your child drinks plenty of fluids and does not become dehydrated. Signs of dehydration include: a dry sticky mouth, decreased urine output (little or no urine output for 8 hours) or dark urine, dizziness or lightheadedness, lethargy or decreased energy or playfulness, or lack of tears. If you think your child is becoming dehydrated, please call your doctor or go to the emergency room.

If your child is sick or has a fever within 24 hours of the time of his/her procedure, please notify our office immediately.

All lab work ordered must be completed and resulted before the procedure to avoid the possibly of rescheduling.

Turkey Creek Medical Center is located at 10820 Parkside Dr, Knoxville TN 37934 and you will park and enter at the smaller building to the left of the main hospital. They will call you the day before the procedure to give you an arrival time. A parent or legal guardian must be present for the entirety of the procedure.

FOLLOW THESE DIRECTIONS **THE DAY BEFORE** YOUR PROCEDURE:

Step One:

Your child may have only clear liquids on the day before the procedure. Clear liquids (you should be able to see light through it) include: Pedialyte, Gatorade, Powerade, light colored juices, tea, clear carbonated drinks, clear broth, bouillon, Jell-O and Popsicles. NO RED fluids should be given. There is no limit to the amount of clear liquids your child may have and we encourage you to offer plenty of fluids to ensure that your child does not become dehydrated.

Step Two:

- At 8:00 AM have your child drink 2 ounces of Magnesium Citrate (over the counter) followed by 6 ounces of Pedialyte, Gatorade, or Powerade.

Step Three:

- At 10:00 AM mix 8 capfuls of Miralax with 16 ounces of Pedialyte, Gatorade, or Powerade. Allow your child to drink this until finished over the next 3 hours. Mixing capfuls into different flavors/beverages may be helpful to your child. Mixing in warm liquid may help it dissolve better.
- 1 tablet of Dulcolax (5mg laxative) with 8 ounces of Gatorade or Powerade.

At 12:00 PM: If your child is not tolerating the prep, please call the office for further instructions.

Step Four:

- At 4:00 PM have your child drink 2 ounces of Magnesium Citrate (over the counter) followed by 6 ounces of Pedialyte, Gatorade or Powerade

Step Five:

- At 7:00 PM have your child take 1 tablet of Dulcolax (5mg laxative) with 8 ounces of Gatorade or Powerade.

At 10:00 PM: If your child still does not have clear watery diarrhea then repeat Step Four.

YOUR CHILD CANNOT HAVE ANYTHING TO EAT OR DRINK AFTER MIDNIGHT UNLESS OTHERWISE DIRECTED BY THE SURGERY DEPARTMENT.